

A journal of world insect systematics

INSECTA MUNDI

0795

New state and island records of Coccinellidae (Coleoptera)
in Hawai'i, USA

Louis S. Hesler

USDA Agricultural Research Service, 2923 Medary Avenue, Brookings, SD 57006, USA
louis.hesler@usda.gov

William D. Perreira

P.O. Box 61547, Honolulu, HI 96839, USA

Dana Anne Yee

Dana Anne Yee Foundation, 1717 Mott Smith Drive #905, Honolulu, HI 96822, USA

Joshua H.S. Silva

University of Hawai'i Cooperative Extension, 955 Kamehameha Highway,
University of Hawai'i at Manoa, Pearl City, HI 96782, USA

Date of issue: September 25, 2020

Center for Systematic Entomology, Inc., Gainesville, FL

Hesler LS, Perreira WD, Yee DA, Silva JHS. 2020. New state and island records of Coccinellidae (Coleoptera) in Hawai'i, USA. *Insecta Mundi* 0795: 1–4.

Published on September 25, 2020 by
Center for Systematic Entomology, Inc.
P.O. Box 141874
Gainesville, FL 32614-1874 USA
<http://centerforsystematicentomology.org/>

INSECTA MUNDI is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. *Insecta Mundi* will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. *Insecta Mundi* publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources, including the Zoological Record and CAB Abstracts. *Insecta Mundi* is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Guidelines and requirements for the preparation of manuscripts are available on the *Insecta Mundi* website at <http://centerforsystematicentomology.org/insectamundi/>

Chief Editor: David Plotkin, insectamundi@gmail.com
Assistant Editor: Paul E. Skelley, insectamundi@gmail.com
Layout Editor: Robert G. Forsyth
Editorial Board: Davide Dal Pos, Oliver Keller, M. J. Paulsen
Founding Editors: Ross H. Arnett, Jr., J. H. Frank, Virendra Gupta, John B. Heppner, Lionel A. Stange, Michael C. Thomas, Robert E. Woodruff
Review Editors: Listed on the *Insecta Mundi* webpage

Printed copies (ISSN 0749-6737) annually deposited in libraries:

| | |
|--|---|
| CSIRO, Canberra, ACT, Australia | Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA |
| Museu de Zoologia, São Paulo, Brazil | Field Museum of Natural History, Chicago, IL, USA |
| Agriculture and Agrifood Canada, Ottawa, ON, Canada | National Museum of Natural History, Smithsonian Institution, Washington, DC, USA |
| The Natural History Museum, London, UK | Zoological Institute of Russian Academy of Sciences, Saint- Petersburg, Russia |
| Muzeum i Instytut Zoologii PAN, Warsaw, Poland | |
| National Taiwan University, Taipei, Taiwan | |
| California Academy of Sciences, San Francisco, CA, USA | |

Electronic copies (online ISSN 1942-1354, CDROM ISSN 1942-1362) in PDF format.

Printed CD or DVD mailed to all members at end of year. Archived digitally by Portico.
Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>
University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>
Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

Copyright held by the author(s). This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited. <http://creativecommons.org/licenses/by-nc/3.0/>

New state and island records of Coccinellidae (Coleoptera) in Hawai‘i, USA

Louis S. Hesler

USDA Agricultural Research Service, 2923 Medary Avenue, Brookings, SD 57006, USA
louis.hesler@usda.gov

William D. Perreira

P.O. Box 61547, Honolulu, HI 96839, USA

Dana Anne Yee

Dana Anne Yee Foundation, 1717 Mott Smith Drive #905, Honolulu, HI 96822, USA

Joshua H.S. Silva

University of Hawai‘i Cooperative Extension, 955 Kamehameha Highway,
University of Hawai‘i at Manoa, Pearl City, HI 96782, USA

Abstract. Three new records of Coccinellidae (Coleoptera) are recorded within the state of Hawai‘i, USA. *Diomus roseicollis* (Mulsant, 1853) is newly recorded from the state based on four specimens collected on the island of O‘ahu. *Cycloneda sanguinea sanguinea* (Linnaeus, 1763) and *Brumoides suturalis* (Fabricius, 1789), both already established in the state, are newly recorded from the island of Hawai‘i. Additional sampling is recommended to determine the distribution of *D. roseicollis* within the island of O‘ahu. Further surveys are warranted to determine particular habitats and prey for *D. roseicollis*, *C. sanguinea sanguinea*, and *B. suturalis* in Hawai‘i, as well as their impact on the insect fauna there.

Key words. Lady beetles, non-indigenous species, new geographic distribution records.

ZooBank registration. urn:lsid:zoobank.org:pub:B5DFA460-E9DB-4146-9FDA-EA04C3CF8DB8

Introduction

The 48 species of lady beetles (Coleoptera: Coccinellidae) on the Hawaiian Islands of the USA have accrued over time from purposeful and accidental introductions since the 1800s (Leeper 1976, 2015; Nishida 2002; Hesler et al. 2019). Purposeful introductions of predators such as lady beetles for biological pest control in Hawai‘i have become generally disfavored and highly scrutinized due to unwitting effects on non-target organisms (Howarth 1985, 2001). However, inadvertent introductions of lady beetles are still possible because of Hawai‘i’s high annual volume of sea and air traffic, and thus additional species may be newly recorded in the future. In addition, the distribution of lady beetles already established in Hawai‘i may be extended because of additional introduction or inter-island movement. Indeed, we report here on a species of lady beetle newly recorded in Hawai‘i and two established species with new island records in the state.

Materials and Methods

Various insects were collected during 2017–2020 as bycatch on yellow sticky board traps (YSBT) used in surveys for assorted invasive plant pests on the islands of O‘ahu and Hawai‘i. Lady beetles were sorted from other insects in the samples and identified using keys in Gordon (1976) and Leeper (2016) and checked against records of lady beetles in Hawai‘i according to Leeper (1976, 2015), Nishida (2002) and Hesler et al. (2019). Label information for specimens was reported verbatim. Pinned specimens of the lady beetles newly recorded in this paper were

labeled as voucher specimens and deposited indefinitely at the USDA-ARS North Central Agricultural Research Laboratory, Brookings, SD, USA. Nomenclature is based on Seago et al. (2011).

Results

***Diomus roseicollis* (Mulsant, 1853) (Coleoptera: Coccinellidae: Coccinellinae: Diomini)**

Hawai'i (**new state record**), Honolulu County, O'ahu Island, Barbers Point, 0–5 ft elev., 27-III to 5-IV-2017, W. D. Perreira and D. A. Yee, YSBT, 1 ♀; Honolulu County, O'ahu Island, Barbers Point, 0–5 ft elev., 5-12-IV-2017, W. D. Perreira and D. A. Yee, YSBT, 1 ♀; Honolulu County, O'ahu Island, Barbers Point, 0–5 ft elev., 28-VI to -2-VIII-2017, W. D. Perreira and D. A. Yee, YSBT, 1 ♀; Honolulu County, O'ahu Island, Waimanalo, UH Ag Res Sta, 70 ft elev., 25-VIII-2017, W. D. Perreira and D. A. Yee, 1 ♀ (Figure 1).

***Brumoides suturalis* (Fabricius, 1789) (Coleoptera: Coccinellidae: Coccinellinae: Chilocorini)**

Hawai'i, Hawai'i County, Hawai'i Island (**new county and new island record**), UH Mealani Res. Sta., 20°02'12"N, 156°36'28"W, el 2880 ft, sweeping, 6-XII-2019, W. D. Perreira, 2 specimens.

***Cycloneda sanguinea sanguinea* (Linnaeus, 1763) (Coleoptera: Coccinellidae: Coccinellinae: Coccinellini)**

Hawai'i, Hawai'i County, Hawai'i Island (**new county and new island record**), 19°47'46"N, 155°37'37"W, Saddle Rd. Rte. 200, 5480 ft, Golden Rod, 22-XI-2019, W. D. Perreira, 3 specimens; 19°47'45"N, 155°37'37"W, Saddle Rd. Rte. 200, el 5500 ft, Golden Rod, 3-I-2020, W. D. Perreira, 1 specimen.



Figure 1. One of four specimens of *Diomus roseicollis* (female) collected from the island of O'ahu constituting a new state record for Hawai'i, USA.

Discussion

Diomus roseicollis joins two other diomine lady beetles, *Diomus debilis* (LeConte) and *Diomus notescens* (Blackburn), already established in Hawai'i (Leeper 2015). The external morphology of *D. roseicollis* includes a blackish-brown elytron with a tan apical spot that does not reach the elytral suture (Gordon 1976) (Figure 1). This characteristic distinguishes *D. roseicollis* from congeneric species, including *D. debilis* and *D. notescens* (Gordon 1976, Leeper 2016).

No prey or habitat data were recorded with the specimens of *D. roseicollis*. However, species of *Diomus* prey upon various kinds of Hemiptera, including Coccoidea and Aphidoidea (Gordon 1999; Hodék and Evans 2012). For instance, *D. roseicollis* is known to prey on the sugarcane aphid, *Melanaphis sacchari* (Zehntner) (Hemiptera: Aphididae) (Rodríguez-Vélez et al. 2019), a pest which is present in Hawai'i (Zimmerman 1948). Surveys should be conducted to determine the extent to which *D. roseicollis* may be preying upon *M. sacchari* and other hemipterans in Hawai'i. Surveys for *D. roseicollis* should be conducted in various habitats and extended to other islands of Hawai'i in order to determine its geographic extent and habitat preferences. The benefits and detriments of its predation on the Hawaiian insect fauna should be assessed.

Brumoides suturalis is native to southern Asia and preys on various insect herbivores, including aphids (Hemiptera: Aphididae) (Swaminathan et al. 2015; Shankar et al. 2015), coccids (Hemiptera: Coccidae) (Gautum 1990), and mealybugs (Hemiptera: Aleyrodidae) (Inayatullah 1984). It had been intentionally introduced into the state of Hawai'i more than once, but apparently established on the islands of Ni'i'hau and O'ahu from accidental introduction (Mau 1976; Leeper 2015). Means of its advent on the island of Hawai'i is unknown, but interisland movement is likely, either by natural dispersal or inadvertent importation by humans. Despite its relatively long period of establishment in the state of Hawai'i, we did not find published prey records for *B. suturalis*. Thus, further surveys are warranted specifically to ascertain its prey in Hawai'i and determine the benefits and detriments of its predation (Howarth 1985).

Cycloneda s. sanguinea is a Nearctic lady beetle (Vandenberg 2002) that has been introduced several times into Hawai'i to control pest aphids (Lai and Funasaki 1986; Leeper 2015). It has been recorded previously from the islands of Maui (Kumashiro et al. 2001) and O'ahu (Hesler et al. 2019). The origin of *C. s. sanguinea* on the island of Hawai'i is unknown, but interisland dispersal is likely, either naturally or facilitated by humans. *Cycloneda s. sanguinea* is principally aphidophagous (Hodék and Evans 2012), but was not associated with any particular prey in our sampling. Follow-up surveys are warranted to determine its prey and the utility and disadvantages of its predation in Hawai'i.

Acknowledgments

The work was funded through USDA-ARS CRIS Project Number 3080-21220-006-00D, The Dana Anne Yee Foundation, The Atherton Family Foundation, Landscape Association of Hawai'i, The O'ahu Nursery Growers Association, and Rick and Whitney Benedict of Beachside Lighting Hawai'i. Special thanks to the following individuals for allowing us to collect on parts of Hawai'i Island: Marla Fergerstrom, University of Hawai'i, Mealani Research Station; Robert Lee, State of Hawai'i, Department of Transportation; and Nahua Guilloz, Parker Ranch Foundation Trust and Parker Ranch, Inc. Eric Beckendorf produced the figure image. Eric Beckendorf, Lauren Hesler, John Leeper and Janakiraman Poorani reviewed drafts of this paper.

Literature Cited

- Gautam RD. 1990.** Mass-multiplication technique of coccinellid predator, ladybird beetle (*Brumoides suturalis*). Indian Journal of Agricultural Sciences 60: 747–750.
- Gordon RD. 1976.** The Scymnini (Coleoptera: Coccinellidae) of the United States and Canada: Key to genera and revision of *Scymnus*, *Nephus*, and *Diomus*. Bulletin of the Buffalo Society of Natural Sciences 28: 1–362.
- Gordon RD. 1999.** South American Coccinellidae (Coleoptera). Part VI. A systematic revision of the South American Diomini, new tribe (Scymninae). Annales Zoologici (Warszawa) 49: 1–219.

- Hesler LS, Ferreira WD, Yee DA, Silva JHS. 2019.** New records of Coccinellidae (Coleoptera) from Hawai'i, U.S.A. *The Coleopterists Bulletin* 73: 382–386.
- Hodék I, Evans EW. 2012.** Food relationships. p. 141–274. *In: Hodék I, van Emden HF, Honěk A. (eds.). Ecology and behaviour of the ladybird beetles (Coccinellidae).* Wiley-Blackwell, John Wiley and Sons; West Sussex, UK. 600 p.
- Howarth FG. 1985.** Impacts of alien land arthropods and mollusks on native plants and animals in Hawaii. p. 149–179. *In: Stone CP, Scott JM (eds.). Hawaii's terrestrial ecosystems: Preservation and management.* Cooperative National Park Resources Studies Unit and University of Hawaii; Honolulu, HI. 584 p.
- Howarth FG. 2001.** Environmental issues concerning the importation of non-indigenous biological control agents. p. 70–91. *In: Lockwood JA, Howarth FG, Purcell MF (eds.). Balancing nature: Assessing the impact of importing non-native biological control agents (An international perspective).* Entomological Society of America; Lanham, MD. 130 p.
- Inayatullah C. 1984.** Sugarcane aleurodids, *Aleurolobus barodensis* (Maskell) and *Neomaskellia andropogonis* Corbett (Hom.: Aleyrodidae), and their natural enemies in Pakistan. *International Journal of Tropical Insect Science* 5: 279–282.
- Kumashiro BR, Heu RA, Nishida GM, Beardsley JW. 2001.** New state records of immigrant insects in the Hawaiian Islands for the year 1999. *Proceedings of the Hawaiian Entomological Society* 35: 171–182.
- Lai PY, Funasaki GY. 1986.** List of beneficial organisms purposely introduced for biological control in Hawai'i: 1890–1986. *Hawai'i State Department of Agriculture Report*; Honolulu, HI. 78 p.
- Leeper JR. 1976.** A review of the Hawaiian Coccinellidae. *Proceedings of the Hawaiian Entomological Society* 22: 279–306.
- Leeper JR. 2015.** An Annotated Checklist of the Coccinellid (Coleoptera: Coccinellidae) Introductions and Establishments in Hawaii: 1885 to 2015. eVols series, University of Hawaii at Manoa; Honolulu, HI. Available at <http://hdl.handle.net/10524/48742>. (Last accessed April 2020.)
- Leeper JR. 2016.** Key to the Coccinellids (Coleoptera: Coccinellidae) Established in Hawaii. eVols series, University of Hawaii at Manoa; Honolulu, HI. Available at <http://hdl.handle.net/10524/49028>. (Last accessed April 2020.)
- Mau R. 1976.** *Brumoides suturalis* (Fabricius). Notes and exhibitions for August 11, 1974 meeting. *Proceedings of the Hawaiian Entomological Society* 22(2): 170.
- Nishida GM. 2002.** Hawaiian terrestrial arthropod checklist. Fourth edition. Bishop Museum technical report 22. Bishop Museum; Honolulu, HI. iv + 310 p.
- Rodríguez-Vélez JM, Gallou A, Uribe-Mú CA, Najar-Pacheco MA, Huerta-Martínez FM, Contreras-Ramos A, Arredondo-Bernal HC. 2019.** Identification of twelve species of Coccinellidae (Coleoptera) predatory on *Melanaphis sacchari* (Zehntner) (Hemiptera: Aphididae) in Mexico, and submission of reference COI sequences. *The Coleopterists Bulletin* 73: 243–251.
- Seago AE, Giorgi JA, Li J, Ślipiński A. 2011.** Phylogeny, classification and evolution of ladybird beetles (Coleoptera: Coccinellidae) based on simultaneous analysis of molecular and morphological data. *Molecular Phylogenetics and Evolution* 60: 137–151.
- Shankar C, Chintagunta L, Muthusamy S, Vailla S, Srinivasan A, Katti G. 2018.** Flora surrounding rice fields as a source of alternative prey for coccinellids feeding on the pests of rice. *European Journal of Entomology* 115: 364–371.
- Swaminathan R, Meena A, Meena BM. 2015.** Diversity and predation potential of major aphidophagous predators in maize. *Applied Ecology and Environmental Research* 13: 1069–1084.
- Vandenberg NJ. 2002.** The New World genus *Cycloneda* Crotch (Coleoptera: Coccinellidae: Coccinellini): Historical review, new diagnosis, new generic and specific synonyms, and an improved key to North American species. *Proceedings of the Entomological Society of Washington* 104: 221–236.
- Zimmerman EC. 1948.** *Insects of Hawaii*. Vol. 5. Homoptera: Sternorrhyncha. University of Hawaii Press; Honolulu, HI. 464 p.

Received June 8, 2020; accepted August 13, 2020.

Review editor Oliver Keller.